

Original Article

Multicultural Sensitivity and Related Factors in Nursing Students of Zanjan University of Medical Sciences During the COVID-19 Pandemic

Zeynab Bakhteri¹ , Nasrin Hanifi² , Nasrin Jafari Varjoshani² 

¹ Master of student, School of Nursing and Midwifery, Zanjan University of Medical Sciences, Iran

² Ph.D. Department in Nursing, Associate Professor, School of Nursing and Midwifery, Zanjan University of Medical Sciences, Iran

Article Info



Article history:

Received 15 October 2021
Accepted 20 June 2022
Published 2 August 2022

Keywords:

Multicultural sensitivity
Cultural Diversity
Students, Nursing
COVID-19

*Corresponding author:

Nasrin Hanif, Department of
Nursing, Associate Professor
School of Nursing and Midwifery
Zanjan University of Medical Sciences
Iran

Email: nasrinhani@zums.ac.ir

Abstract

Background & Objective: Holistic care is provided by the ability to communicate effectively with patients from different cultures. In this respect, multicultural sensitivity helps communicate effectively with the patient. Therefore, the present study was conducted to investigate the multicultural sensitivity of nursing students and the factors affecting it during the COVID-19 pandemic.

Materials & Methods: This cross-sectional correlation study was conducted on 245 nursing students of the Zanjan University of Medical Sciences in 2021. Data were collected using two questionnaires of demographic information and multicultural sensitivity. The multicultural questionnaire scored multicultural sensitivity on a Likert scale in the range of 1 to 5. Scores ≤ 3 were unfavorable multicultural sensitivities > 3 were desirable cultural sensitivities. Data analysis was performed using an independent t-test, analysis of variance, and logistic regression in the SPSS statistical software.

Results: The mean total score of multicultural sensitivity was 3.64 ± 0.48 . Also, the mean of multicultural sensitivity based on gender, residence, religion, bilingualism, semester, and field of study did not show a statistically significant difference ($P < 0.05$). The logistic regression model showed that gender and field of study could predict 3 to 6% of changes in multicultural sensitivity.

Conclusion: The findings of this study showed that the multicultural sensitivity of students is favorable, despite their distance from the university environment. However, it is recommended to conduct training to make them more familiar with the cultural habits and health of Iranian ethnic groups.



Copyright © 2021, This is an original open-access article distributed under the terms of the Creative Commons Attribution-noncommercial 4.0 International License which permit copy and redistribution of the material just in noncommercial usages with proper citation

Introduction

Cultural diversity means differences in lifestyle, language, values, norms, and other components of culture within or between groups (1). This diversity is increasing in different communities because of job opportunities, better living conditions, and access to health facilities and other facilities (2, 3). Different cultures pose many challenges to communication in clinical settings and can lead to adverse outcomes for patients (4). Nurses have to communicate effectively with patients from different cultures to meet the needs of their patients and prevent adverse outcomes (5). Therefore, all physical, mental, emotional, spiritual, social, and economic aspects of the patients are considered with an emphasis on their dignity.

Holistic care in nursing is comprehensive patient care in which the nurse's knowledge, theories, specialties, insights, and creativity are used. Therefore, the patients' physical, mental, emotional, spiritual, social, and economic aspects are considered with emphasis on their dignity (6). Holistic care occurs when the nurse understands the patient's cultural values and is aware of

and sensitive to the cultural differences between oneself and the patient (7).

Multicultural sensitivity refers to the ability to respond appropriately to the attitudes, feelings, or circumstances of different races, nationalities, religions, languages, and cultures (1). Multicultural sensitivity is a dimension of cultural competence by which one can respond appropriately to the attitudes, feelings, or circumstances of people with different racial, religious, linguistic, and cultural heritage (8). Tucker et al. introduced patient-centered care with multicultural sensitivity and emphasized paying attention to patient needs and wants and patient empowerment. Such care makes the patient feel comfortable, respected, and trusted in service delivery (9).

In (3, 10), the multicultural sensitivity of nurses was moderate. However, in (11), the multicultural sensitivity of nurses in suburban hospitals exposed to diverse cultures was reported to be high. A positive association between multicultural sensitivity and assertiveness has also been reported (10).

Valizadeh et al. showed that culturally sensitive care in caring for a sick child has consequences such as honesty of parents in providing information and expression of cultural needs, family satisfaction, reducing family stress, promoting nurse-child/family communication, and preventing misunderstandings (1).

Iran is a multicultural country, and ethnic and linguistic diversity is one of its key features. Because of its privileged position as a global crossroads, the historical land of Iran has always been the crossroads of different cultures and civilizations and the arena of various movements and invasions (12). The coexistence of Persian, Azeri, Kurdish, Gilak, Lurs, Arab, Mazani, Baluch, Bakhtiari, Talesh, and Turkmen ethnic groups causes cultural diversity in this country. Therefore, different religions and ethnicities in a community raise the need to pay attention to culture in the face of individuals (1). Zanjan is an Iranian immigrant city whose urban population has increased from 28.8% to 63.4% during the last 40 years (13). This increase in Zanjan's population has created great cultural diversity in the city. Most medical and paramedical students, especially nursing students, are non-native and do not speak Turkish. Most illiterate patients speak only their local language and dialect, which complicates the communication between students and patients. Besides, patient and family dissatisfaction affects therapeutic interventions and optimal care. Zanjan province is composed of the Aryan, Georgian, and Khamseh tribes. The people of this province speak Azeri Turkish, but the proximity to the capital (Tehran) has led to a tendency toward Persian. In addition, Turkish is mixed with Gilaki in its north, Kurdish in its southwest, Azerbaijani Turkish in its west, and Tati in its south (13). Multicultural sensitivity is made possible by the ability to communicate effectively with patients from different cultures (5). In the COVID-19 pandemic, where the presence of students in clinical fields has diminished, students' multicultural sensitivity decreased due to less interaction with patients. Communication with patients affected and prevented face-to-face communication and receiving non-verbal messages due to using personal protective equipment such as masks and face shields, interaction with patients, and increased multicultural sensitivity in students. This problem was more common in nursing and other medical students who were non-native and had a distinct language and dialect from hospitalized patients. Evaluating the development of multicultural sensitivity and related factors in nursing students is essential during their studies. Therefore, regarding the knowledge gap in the status of multicultural sensitivity and related factors in nursing students of the Zanjan University of Medical Sciences, especially in pandemic

COVID-19, this study aims to determine the level of multicultural sensitivity and related factors in nursing students of the Zanjan University of Medical Sciences in COVID-19 pandemic.

Material & Methods

This was a descriptive cross-sectional study. The study population included students of the Faculty of Nursing affiliated with the Zanjan University of Medical Sciences. The Zanjan University of Medical Sciences comprises eight faculties. In the School of Nursing, 600 undergraduate students are studying in the fields of nursing, midwifery, operating room nursing, and anesthesia, and 100 students are studying in 5 postgraduate fields. The sample size was estimated by performing a pilot study on 30 students. By considering $\alpha = 0.05$, $\delta = 0.76$, and $d = 0.1$, the sample size was estimated to be 221 as follows:

$$n = \left(\frac{z/\delta}{d} \right)^2$$

Considering a 10% dropout rate, the final sample size was 245 people. Inclusion criteria were willingness to participate in the study, studying in the third semester and above, and studying in one discipline of nursing, midwifery, operating room, and anesthesia. Because the 3rd semester and older students have experience in the clinical setting, this criterion was considered for selecting the participants.

Data collection tools included a demographic information questionnaire and a multicultural sensitivity questionnaire.

The questionnaire included questions about age, gender, semester, place of residence, bilingualism, place of residence, and religion.

The Multicultural Sensitivity Questionnaire was designed by Jibaja et al. (2013). This questionnaire comprises five dimensions and 21 items on a 5-point Likert scale ranging from 1 "strongly disagree" to 5 "strongly agree". Scores ≤ 3 were considered unfavorable multicultural sensitivities and > 3 as desirable multicultural sensitivities. This questionnaire has 5 dimensions: 1) interaction engagement, 2) respect for multicultural differences, 3) interaction confidence, 4) interaction enjoyment, and 5) interaction attentiveness. The reliability of the questionnaire with Cronbach's alpha was reported to be 0.89 (14).

Regarding the validity of the translation process, it was translated based on the standards recommended in the guidelines (15). In this study, two people translated the questionnaire into Persian. Three experts approved the translation of the questionnaire. The content validity index (CVI), based on the opinion of 10 experts, was calculated to be 93% for all items. Then, the

questionnaire was translated into English again and matched with the original questionnaire. Two nursing professionals confirmed the English fluency of the retranslated questionnaire. The reliability of the questionnaire was determined using Cronbach's alpha coefficient of 0.86.

Sampling was done online. The link to the questionnaire and a letter explaining the research objectives and getting informed consent were provided to students through WhatsApp, Telegram, and email. SPSS statistical software version 22 was used for data analysis. After examination by the Kolmogorov-Smirnov test, the normality of the data distribution was determined. Descriptive statistics, frequency, mean, and standard deviation (\pm SD) were used to evaluate the baseline data. In addition, independent t-test, analysis of variance (ANOVA), and post-hoc test with the least significant difference of LSD were used to evaluate the level of multicultural sensitivity based on demographic variables. A logistic regression test was used to predict the predictability of demographic variables in multicultural sensitivity. The significance level was considered less than 0.05.

The ethics committee of Zanjan University of Medical Sciences approved the study with the ethics code IR.ZUMS.REC.1399.201. At the beginning of the online questionnaire, the research objectives were

explained on the first page. If the participants were willing to participate in the study, they were asked to complete the questionnaires by approving the consent form.

Results

A total of 245 nurses were recruited via the online data gathering and their data were analyzed. The results showed that 28.6% (n = 70) of the participants in semester 3, 20.8% (n = 51) in semester 4, 13.9% (n = 33) in semester 5, 21.2% (n = 52) in semester 6, 8.2% in semester 7 (n = 20), and 7.8% (n = 19) were studying in the 8th semester; 63.6% (n = 131) were natives and 46.6% (n = 114) were non-natives; 95.5% (n = 234) were Shiites and 4.5% (n = 11) were Sunnis; 77.2% (n = 189) were bilingual and 22.9% (n = 56) were monolingual.

According to the obtained total multicultural sensitivity score, 12.7% (n = 31) of the participants had unfavorable multicultural sensitivity and 86.9% (n = 213) had favorable multicultural sensitivity. Also, the mean overall score of multicultural sensitivity (3.65 ± 0.48) was at a favorable level. Respect for different cultures had the highest mean (3.96 ± 0.75) interaction confidence had the lowest mean (3.4 ± 0.57) (Table 1).

Table 1. The mean scores of dimensions and the overall score of multicultural sensitivity in participants

| Items | Minimum | Maximum | Mean \pm SD |
|---|---------|---------|-----------------|
| Interaction engagement | 1.57 | 4.9 | 3.5 ± 0.52 |
| Respect of cultural differences | 1.17 | 5 | 3.96 ± 0.75 |
| Interaction confidence | 1.6 | 4.8 | 3.4 ± 0.57 |
| Interaction enjoyment | 1 | 5 | 3.8 ± 0.87 |
| Interaction attentiveness | 1 | 5 | 3.6 ± 0.62 |
| The overall score of cultural sensitivity | 2.19 | 4.67 | 3.65 ± 0.48 |

SD = Standard Deviation

The results of the independent t-test did not show a statistically significant difference between the mean overall score of multicultural sensitivity based on gender, residence, bilingualism, and religion ($P < 0.05$) (Table 2).

ANOVA results for the difference in the mean overall score of multicultural sensitivity based on the variables of the field of study and the semester did not show a significant difference (Table 3).

Table 2. Comparison of the mean overall score of participants' multicultural sensitivity based on the variables of gender, residence, religion, and bilingualism

| Variable | T-independent test | | | |
|------------------------|--------------------|-------------|--------|-----|
| | Mean ± SD | t | df | P |
| Gender | Men | 3.69 ± 0.54 | -1.28 | 243 |
| | Women | 3.67 ± 0.44 | | |
| Residence | Native | 3.6 ± 0.47 | -1.34 | 243 |
| | Non-native | 3.7 ± 0.49 | | |
| Religion | Shia | 3.6 ± 0.49 | -1.39 | 243 |
| | Sunni | 3.8 ± 0.37 | | |
| Being bilingual | Yes | 3.63 ± 0.49 | -0.707 | 243 |
| | No | 3.96 ± 0.45 | | |

SD = Standard Deviation

Table 3. Comparison of the mean overall score of participants' multicultural sensitivity based on the majors and semester variables

| Variable | ANOVA | | | |
|-----------------|------------------------|-------------|------|---|
| | Mean ± SD | F | df | P |
| Majors | Nursing | 3.66 ± 0.46 | 1.67 | 3 |
| | Operating room nursing | 3.68 ± 0.56 | | |
| | Anesthesia nursing | 3.75 ± 0.54 | | |
| | Midwifery | 3.47 ± 0.45 | | |
| Semester | Three | 3.6 ± 0.53 | 2.16 | 5 |
| | Four | 3.7 ± 0.47 | | |
| | Five | 3.7 ± 0.49 | | |
| | Six | 3.5 ± 0.41 | | |
| | Seven | 3.5 ± 0.43 | | |
| | Eight | 3.7 ± 0.47 | | |

SD; Standard Deviation

A logistic regression model with the forward likelihood ratio method was used to investigate the predictors of multicultural sensitivity. The model was implemented in two steps. The results of the omnibus holistic test showed that the model fits in the second step (chi-square = 67.8, df = 1). The results showed that two variables of gender and majors could predict students' cultural sensitivity ($P < 0.013$).

The Cox & Snell R-square and Nagelkerke R-square values were 0.035 and 0.067, respectively. These coefficients showed that these two variables could only predict 3 to 6% of multicultural sensitivity changes. The odds ratios for gender and majors were 3.13 and 0.66, respectively. Therefore, gender and major increased multicultural sensitivity by 3.13 and 0.66 times (Table 4).

Table 4. Logistic regression analysis to predict multicultural sensitivity in the participants

| Variable | Logistic regression analysis | | | | | |
|----------|------------------------------|-------|------|----|-------|--------|
| | B | S.E | Wald | df | P | EXP(B) |
| constant | 0.89 | 0.64 | 1.93 | 1 | 0.16 | 2.42 |
| Gender | 1.14 | 0.448 | 6.46 | 1 | 0.011 | 3.13 |
| Majors | -0.422 | 0.19 | 4.99 | 1 | 0.025 | 0.66 |

Discussion

This study's result showed that the multicultural sensitivity of students was in favorable conditions. The mean of multicultural sensitivity based on gender, residence, religion, bilingualism, semester, and majors did not show a statistically significant difference. Also, the logistic regression model results showed that gender and majors could predict only 3 to 6% of multicultural sensitivity changes.

In a study in Turkey, Yilmaz et al. (2017) reported that the multicultural sensitivity of nurses in suburban hospitals promoted multicultural sensitivity (11) because of their greater contact with patients from different cultures. Also, the study by Zialeme et al. (2016) in Tabriz (Iran) showed that the mean score of multicultural sensitivity of nurses was moderate (3). This difference can be related to nurses' training in knowing different cultures. Also, another study showed that the multicultural sensitivity of nursing students was moderate (10). Therefore, the relationship of the students of the present study with patients from different cultures has promoted their multicultural sensitivity.

Also, the results showed that the highest mean score of multicultural sensitivity was related to the dimension of respect for cultural differences, and the lowest mean was related to the dimension of interaction confidence. The high score on the dimension of respect for cultural differences indicates a high understanding of students of the cultural differences between them and patients.

Low interaction confidence scores can be due to students' inability to communicate effectively due to the COVID-19 pandemic conditions and using personal protective equipment (e.g., masks, shields, and glasses), fear of misunderstandings due to cultural differences, and fear of the unknown. This finding can also be related to the lack of essential intercultural education in educating students in both studies and, therefore, the lack of confidence in interaction with patients from different cultures. Teaching how to communicate, teaching the customs of different cultures, and helping resolve cultural conflicts can increase students' confidence in interaction. In the study by Toda et al. (2018) in Japan, the lowest score was related to the dimension of interaction confidence (5). In another

study by Erkin et al. (2017) in Turkey, the highest and lowest mean of multicultural sensitivity were related to the dimensions of interaction engagement and interaction enjoyment, respectively (16).

Also, according to the results of this study, gender and majors were determined as predictors of multicultural sensitivity. Therefore, women were more culturally sensitive than men. Kılıç et al. reported a significant relationship between gender and cultural sensitivity (10). The similarity of the findings could be related to the high proportion of women participants in both studies. One of the effective factors is the existence of attitudes rooted in stereotyped thoughts about each ethnic group that manifest themselves in the value judgments of human beings (17). A strategy to promote multicultural sensitivity is the tendency toward a cultural approach to the ethnic phenomenon. This approach considers customs, subcultures, and all cultural components related to ethnic groups, local cultures, and ethnicities (18). Also, improving the cultural competence of nurses increases the observance of patients' rights, attention to their cultural values, and efforts to reduce patients' suffering (19). Therefore, strategies should be taken to increase the multicultural sensitivity of students, including familiarity with the values governing different cultures and ethnicities.

Conclusion

The present study shows the need to pay attention to the multicultural sensitivity of medical sciences students. Respect for the cultures, values, and beliefs of individuals in the community is a professional responsibility and is in line with the autonomy of individuals. Given the importance of the impact of culture on people's health and its role in community health, those in charge of medical education should familiarize medical students with existing subcultures and be effective in health. However, promoting multicultural sensitivity, familiarizing students with different cultures in the community, and accepting and respecting them should be done from the years before students enter the university to institutionalize respect for different cultures. Overall, it is suggested to conduct

research in this field to know more about these cultures. The explanation is that better knowledge of subcultures by the medical staff causes effective interventions to strengthen the correct beliefs about people's health.

Limitation

One limitation of this study was using an electronic questionnaire because of the COVID-19 pandemic conditions, which could limit the generalizability of the findings. Therefore, using probabilistic sampling is recommended for research in a larger statistical population.

Acknowledgements

This article results from a research project by the Faculty of Nursing and Midwifery of the Zanjan University of Medical Sciences and the Vice Chancellor for Research with the code IR.ZUMS.REC.1399.201. The authors would like to thank the Vice Chancellor for the Research of the Zanjan University of Medical Sciences and all the students of the School of Nursing and Midwifery who helped us in this research.

Conflicts of interest

The authors have expressed no conflict of interest.

References

1. Valizadeh L, Zamanzadeh V, Ghahramanian A, Aghajari P. Exploring the consequences of culturally sensitive care in pediatric nursing: A qualitative study. *Hayat*. 2017;23(1):59-73. <https://doi.org/10.22038/ijp.2016.7975>.
2. badriazarin y, hooshang t. The Relationship between Cultural Intelligence and Individual Performance in Employees of Sport and Youth General Office of East Azerbaijan Province. *Journal of Sport Management*. 2017;9(1):55-70. <https://doi.org/10.22059/jsm.2017.62274>.
3. Zialame L. Intercultural sensitivity and its relation to professional interaction among the nurses working in Tabriz medical education institutes in 2014: Tabriz University of Medical Sciences, School of Nursing and Midwifery, Aras ...; 2016.
4. Foronda C, MacWilliams B, McArthur E. Interprofessional communication in healthcare: An integrative review. *Nurse education in practice*. 2016;19:36-40. <https://doi.org/10.1016/j.nep.2016.04.005>.
5. Toda T, Maru M. Cultural sensitivity of Japanese nurses: Exploring clinical application of the Intercultural Sensitivity Scale. *Open Journal of Nursing*. 2018;8(9):640-55. <http://doi.org/10.4236/ojn.2018.89048>.
6. Ghorbani F, Salsali M. Concept of Holistic in Nursing: A review article. *Education and Ethics in Nursing*. 2018;7(3):23-30. <https://doi.org/10.52547/ethicnurs.7.3.4.23>.
7. Congleton A. Providing healthcare to the isolated and unreached: the importance of a community needs assessment to build cultural intelligence. MSc dissertation in Nursing. 2019; Liberty University:83-8.
8. Benito LT, Casas J, O'Donohue WT. Training culturally competent psychologists: A systematic review of the training outcome literature. *Training and Education in Professional Psychology*. 2018;12(3):125-34. <https://doi.org/10.1037/tep0000190>.
9. Tucker CM, Arthur TM, Roncoroni J, Wall W, Sanchez J. Patient-Centered, Culturally Sensitive Health Care. *American Journal of Lifestyle Medicine*. 2013;9(1):63-77. <https://doi.org/10.1177/1559827613498065>.
10. Kılıç SP, Sevinç S. The Relationship Between Cultural Sensitivity and Assertiveness in Nursing Students from Turkey. *Journal of transcultural nursing: official journal of the Transcultural Nursing Society*. 2018;29(4):379-86. <https://doi.org/10.1177/1043659617716518>.
11. Yilmaz M, Toksoy S, Direk ZD, Bezirgan S, Boylu M. Cultural Sensitivity Among Clinical Nurses: A Descriptive Study. *Journal of nursing scholarship: an official publication of Sigma Theta Tau International Honor Society of Nursing*. 2017;49(2):153-61. <https://doi.org/10.1111/jnu.12276>.
12. Montazeri A, Baseri B, Faghih Habibi A, Behnia M. Challenges and Opportunities of Cultural Diversity and Human Rights. *International Journal of Political Science*. 2020;10(2):105-24. <https://doi.org/10.21859/priw-090409>.
13. Plan and Budget Organization of the Islamic Republic of Iran (PBO): <https://www.amar.org.ir/english>; 2020 [cited 2020 28 March]. Available from: <https://www.amar.org.ir/english.2020>.
14. Jibaja-Ruth ML, Kingery PM, Holcomb JD, Buckner WP, Pruitt BE. Development of a Multicultural Sensitivity Scale. *Journal of Health Education*. 2013;25(6):350-7. <https://doi.org/10.1080/10556699.1994.10603060>.
15. Beaton D, Bombardier C, Guillemin F, Ferraz MB. Recommendations for the cross-cultural adaptation of health status measures. New York: American Academy of Orthopaedic Surgeons. 2002; 12:1-29.
16. Erkin O, Bayik Temel A. Cultural sensitivity and related factors among nurse educators in Turkey. *International Journal of Caring Sciences*. 2017;10(3):1374-81.
17. Bahrami Paveh R. Analysis of the Development Paradigm and Its Impact on National Convergence in Iran. *Quarterly journal of The Macro and Strategic Policies*. 2018;6(23):478-97. <https://doi.org/10.32598/jmsp.6.3.478>.

18. Fathi Vajargah K, Bazdar ghamcheh ghiyeh M. A Reflection on the Types of Approaches to Ethnicity in Iran and its Effects on the Educational System. *Journal of Management and Planning in Educational System*. 2020;13(1):37-56. <https://doi.org/10.29252/mpes.13.1.37>.

19. Cruz JP, Alquwez N, Cruz CP, Felicilda-Reynaldo RFD, Vitorino LM, Islam SMS. Cultural competence among nursing students in Saudi Arabia: a cross-sectional study. *International nursing review*. 2017; 64 (2): 215-23. <https://doi.org/10.1111/inr.12370>.

Zeynab Bakhteri, Nasrin Hanifi, Nasrin Jafari Varjoshani. Multicultural Sensitivity and Related Factors in Nursing Students of Zanjan University of Medical Sciences during the COVID-19 Pandemic. *J Med Educ Dev*. 2022; 15 (45): 19-25